

TEST YOUR KNOWLEDGE

Test yourself on what you know about anticoagulation therapy.

INSTRUCTIONS: Please select "True" if the statement is true, select "False" if the statement is false

- | | | |
|---|------|-------|
| 1. Coumadin (warfarin) reduces blood clot formation. | True | False |
| 2. A major side effect of too much Coumadin is bleeding. | True | False |
| 3. Foods high in vitamin K can effect blood clotting. | True | False |
| 4. You should leave the Tenderlett [®] Plus on the cuvette all the time after starting the test. | True | False |
| 5. Once opened, the cuvette is good for 14 days. | True | False |
| 6. Results are reported in PT seconds only. | True | False |
| 7. You can use the ProTime test cuvette immediately upon removal from the refrigerator. | True | False |
| 8. You have 10 seconds in which to remove the Tenderlett Plus from the ProTime cuvette after the blood is drawn into the cuvette. | True | False |
| 9. You should read the ProTime Operator's Manual before performing a test. | True | False |
| 10. Blood collection must be completed within 2 minutes and 10 seconds. | True | False |
| 11. Wipe away the first trace of blood before collecting in the sample cup. | True | False |
| 12. The thromboplastin reagent used has an ISI sensitivity near 2.0. | True | False |

Name: _____

Facility: _____

Date: _____

ProTime and Tenderlett are registered trademarks of ITC.



8 Olsen Avenue • Edison, NJ 08820 USA
tel: 732.548.5700 • fax: 732.248.1928
www.itcmed.com

a subsidiary of Thoratec Corporation